

# Prefrontal cortex glutamate and extraversion

Simone Grimm,<sup>1,2,3</sup> Florian Schubert,<sup>4</sup> Maren Jaedke,<sup>5</sup> Jürgen Gallinat,<sup>6</sup> and Malek Bajbouj<sup>1,2,7</sup>

<sup>1</sup>Cluster of Excellence 'Languages of Emotion', <sup>2</sup>Dahlem Institute for Neuroimaging of Emotion, Freie Universität Berlin, Habelschwerdter Allee 45, 14195 Berlin, Germany, <sup>3</sup>Department of Psychiatry, University Zurich, Lenggstrasse 31, 8032 Zurich, Switzerland,

<sup>4</sup>Physikalisch-Technische Bundesanstalt, (PTB), Abbestr. 2-12, 10587 Berlin, Germany, <sup>5</sup>Department of Psychiatry, Jewish Hospital Berlin,

<sup>6</sup>Department of Psychiatry, Charité, CCM, Charitéplatz 1, 10117 Berlin and <sup>7</sup>Department of Psychiatry, Charité, CBF, Eschenallee 3, 14050 Berlin, Germany

**Extraversion is considered one of the core traits of personality. Low extraversion has been associated with increased vulnerability to affective and anxiety disorders. Brain imaging studies have linked extraversion, approach behaviour and the production of positive emotional states to the dorsolateral prefrontal cortex (DLPFC) and glutamatergic neurotransmission. However, the relationship between extraversion and glutamate in the DLPFC has not been investigated so far. In order to address this issue, absolute glutamate concentrations in the DLPFC and the visual cortex as a control region were measured by 3-Tesla proton magnetic resonance spectroscopy (1H-MRS) in 29 subjects with high and low extraversion. We found increased glutamate levels in the DLPFC of introverts as compared with extraverts. The increased glutamate concentration was specific for the DLPFC and negatively associated with state anxiety. Although preliminary, results indicate altered top-down control of DLPFC due to reduced glutamate concentration as a function of extraversion. Glutamate measurement with 1H-MRS may facilitate the understanding of biological underpinnings of personality traits and psychiatric diseases associated with dysfunctions in approach behaviour and the production of positive emotional states.**

**Keywords:** 1H-MRS; glutamate; prefrontal cortex; personality; extraversion

## INTRODUCTION

The neuronal signatures of individual differences in human personality traits are of growing interest for neuropsychological and psychiatric research. Personality refers to an integrated pattern of thinking, feeling and behaving that varies among individuals but is stable within each individual over time (Sugiura *et al.*, 2000). According to the five-factor model (FEM) of personality, the dominant model of normal personality (Livesley, 2001), all phenotypical variation present in overt behaviour can be explained by five higher order factors comprising extraversion, neuroticism, openness, agreeableness and conscientiousness (Goldberg, 1992). Extraversion and neuroticism are the two core traits of personality that have achieved general acceptance, whereas the importance and definition of the other three factors are still being discussed (Depue and Lenzenweger, 2001). Individuals high on the extraversion dimension are often upbeat, optimistic, enjoy social contact, report more positive emotions in everyday life (McCrae and Costa, 2003) and show greater sensitivity to positive or rewarding stimuli (Lucas and Diener 2001; Banich *et al.*, 2009). Models that attempt to describe biological bases of personality propose lower prefrontal activity in extraverts compared to introverts reflecting less frontally based cognition, such as remembering events from the past, making plans for the future or problem

solving (Gray, 1972; Eysenck, 1986). Behavioural differences in extraverts therefore result from an innate drive to compensate for underactive reticulo-thalamo-cortical pathways (Eysenck, 1986) or lower activity in the behavioural inhibition system, a functional loop that includes the ascending reticular activating system, the frontal lobes, septal regions and the hippocampus (Gray, 1972). Accordingly, early neuroimaging studies that explored the neurobiological underpinnings of extraversion reported a decreased overall cerebral blood flow (Mathew *et al.*, 1984; Stenberg *et al.*, 1990) in extraverts. Johnson *et al.* (1999) specified these findings by showing that extraversion was correlated with enhanced blood flow in the anterior cingulate gyrus, right insular cortex and bilateral temporal lobes, but bilaterally decreased blood flow in the frontal lobe. Other brain regions that have been associated with extraversion in cerebral blood flow studies include the orbitofrontal cortex (Deckersbach *et al.*, 2006) and amygdala (Vaidya *et al.* 2007). Functional imaging studies that explored the neurobiological underpinnings of extraversion have found associations with activity in the amygdala, anterior cingulate cortex, dorsolateral and ventromedial prefrontal cortex and subcortical brain areas (Canli *et al.* 2001, 2002, 2004; Kumari *et al.* 2004; Simon *et al.* 2010; Brühl *et al.*, 2011). Structural studies showed a positive association between extraversion and grey matter concentration for the left amygdala (Omura *et al.*, 2005) as well as a thinner cortical grey matter ribbon in the lateral prefrontal cortex of extraverts (Wright *et al.*, 2006). These findings emphasize the role of the lateral prefrontal cortex

Received 20 June 2011; Accepted 15 August 2011

Advance Access publication 20 October 2011

Correspondence should be addressed to Simone Grimm, PhD, Cluster of Excellence 'Languages of Emotion', Freie Universität Berlin, Habelschwerdter Allee 45, 14195 Berlin, Germany.

E-mail: simone.grimm@fu-berlin.de

for extraversion. The dorsolateral prefrontal cortex (DLPFC) maintains widespread interconnections within the inhibitory top-down control network and is involved in the strategic implementation of control over one's thoughts and actions in line with specific goals or task-oriented behaviours (MacDonald *et al.*, 2000; Botvinick *et al.*, 2001). Several functional imaging studies have reported DLPFC activations during cognitive tasks that require response inhibition (Kelly *et al.*, 2004; Lavric *et al.*, 2004; Barber and Carter, 2005). Dysfunction of the DLPFC also appears to be a key feature of depression, and it has been suggested that dysfunction of the left DLPFC, in particular, may be related to deficits in the ability to experience positive affective states (Davidson *et al.*, 2003; Grimm *et al.*, 2008), which might explain why low trait levels of extraversion have been linked to increased vulnerability to affective and anxiety disorders (Watson and Clark, 1997; Kotov *et al.*, 2010). These disorders, which are marked by specific alterations in the production of positive emotional states and approach behaviour, seem to be associated with glutamatergic neurotransmission (Mathew *et al.*, 2008; Sanacora *et al.*, 2008; Walter *et al.*, 2009; Hashimoto *et al.*, 2010). Despite the links between personality traits and a predisposition towards psychiatric disorders as well as between specifically mood disorders and glutamatergic dysfunction, only few studies have attempted to investigate the role of glutamate for different personality traits. Gallinat *et al.* (2007) could show a negative association between anterior cingulate cortex (ACC) and hippocampal glutamate concentration and sensation seeking. Harm avoidance has also been reported to correlate negatively with glutamate concentration, but positively with GABA concentration in ACC (Kim *et al.*, 2009). Hoerst *et al.* (2010) investigated the association between glutamate levels in ACC with self-reported impulsivity in patients with borderline personality disorder as well as in healthy controls and found a positive correlation irrespective of diagnosis. Another MRS study by Montag *et al.* (2008) found a negative correlation between glutamate concentration in DLPFC and mental perspective taking. To the best of our knowledge, only one study has investigated the association between extraversion and neurotransmitter concentrations (Goto *et al.*, 2010) and reported a negative correlation with GABA in the frontal lobes. All these studies have been limited to correlations between neurotransmitters and personality traits, while no actual differences in neurotransmitter concentrations between subjects scoring high or low on extraversion or other personality dimensions have been reported. Furthermore, the relationship between extraversion and glutamate in the left DLPFC, a key region in the neural network subserving cognitive control, goal-directed behaviour and positive affective states (Davidson *et al.*, 2002), has not been investigated so far. This study aimed to investigate DLPFC glutamate concentration in introverts and extraverts, since several lines of evidence indicate a close relationship between DLPFC glutamate concentration and extraversion.

First, extraverts are psychologically characterized by an increased experience of positive emotions and high approach behaviour (McCrae and Costa, 2003). Secondly, extraverts are neurally characterized by a decreased bilateral cerebral blood flow (Mathew *et al.*, 1984; Stenberg *et al.*, 1990; Johnson *et al.*, 1999) and reduced lateral prefrontal grey matter (Wright *et al.*, 2006). Thirdly, low trait levels of extraversion have been linked to increased vulnerability to major depression and anxiety (Watson and Clark, 1997; Kotov *et al.*, 2010). Lastly, glutamatergic dysfunction has been reported in depression and anxiety disorders and these disorders are also characterized by specific alterations in the production of positive emotional states and approach behaviour (Mathew *et al.*, 2008; Sanacora *et al.*, 2008; Walter *et al.*, 2009; Hashimoto *et al.*, 2010). We hypothesized a lower DLPFC glutamate concentration in extraverts, based on models proposing lower prefrontal activity in extraverts (Gray, 1972; Eysenck, 1986) as well as on previous findings of decreased bilateral cerebral blood flow (Johnson *et al.*, 1999) and reduced gray matter in lateral prefrontal cortex of extraverts (Wright *et al.*, 2006). A further goal was to determine the regional specificity of the hypothesized differences in glutamate concentration between introverts and extraverts by investigating glutamate concentrations in a control region, i.e. the visual cortex (VC). We hypothesized a region-specific difference in DLPFC glutamate concentration.

## MATERIAL AND METHODS

### Subjects

The study was approved by the ethics committee of the Charité University Medicine (Berlin, Germany). Twenty-nine healthy female subjects (age  $32.3 \pm 13.3$  years) were recruited through newspaper advertisements. All subjects gave written informed consent. Somatic as well as psychiatric health status was evaluated by a structured psychiatric interview (Mini-International Neuropsychiatric Interview) (Sheehan *et al.*, 1998) performed by a psychiatrist. No subject had to be excluded due to fulfilling the criteria for an axis I or axis II disorder according to DSM-IV criteria, diagnosed neurological and general medical disorders or clinically relevant abnormalities. Subjects were investigated with magnetic resonance spectroscopy, the NEO five-factor inventory (NEO-FFI, German version, Costa and McCrae, 1989; Borkenau and Ostendorf, 2008) and the State-Trait Anxiety Inventory (STAI; Laux *et al.*, 1981). The NEO-FFI operationalizes the FFM of personality (McCrae and Costa, 1997) and provides domain scores that correspond to the five orthogonal factors extraversion, neuroticism, openness-to-experience, agreeableness and conscientiousness. It consists of 60 self-report items answered on a five-point Likert format. The personality traits are characterized as such (i) extraversion is characterized by talkativeness, assertiveness and energy. (ii) Neuroticism is characterized by upset-ability and is the polar opposite of emotional stability.


(iii) Openness emphasizes originality, curiosity and ingenuity. (iv) Agreeableness is characterized by good-naturedness, cooperativeness and trust. (v) Conscientiousness is characterized by orderliness, responsibility and dependability. The five factors have been shown to be present in psychiatric as well as normative samples (Bagby and Ryder, 2000), and stability estimates for the domains and facets are substantial in both nonclinical (Costa and McCrae, 1992) and patient cohorts (Santor *et al.*, 1997). In addition, evidence from twin studies indicates that each of the five broad trait domains have substantial and unique heritability coefficients (Jang *et al.*, 1998).

The STAI is a self-report assessment device, which includes separate measures of state and trait anxiety. State anxiety reflects a transitory emotional state, may fluctuate over time and can vary in intensity. In contrast, trait anxiety refers to a general tendency to respond with anxiety to perceived threats in the environment.

### MRS procedure

MRS was carried out on a 3-Tesla scanner (MEDSPEC 30/100, Bruker Biospin, Ettlingen, Germany). T1-weighted images were acquired using MDEFT [TE = 3.8 ms, TR = 20.53 ms; 128 contiguous slices, 1.5-mm thick; 1-mm in-plane ( $x-y$ ) resolution]. After localized shimming magnetic resonance spectra were recorded from voxels including the left DLFPFC and the VC (Figure 1). The left DLFPFC voxel

(Brodmann areas 9, 10 and 46) extended  $2 \times 2 \times 2$  cm and was positioned using coordinates as described by Rajkowska and Goldman Rakic (1995). The VC voxel extended  $4 \times 2 \times 2$  cm and was positioned with the anterior border abutting the parietooccipital fissure and centred on the longitudinal fissure. Consistency between subjects was maintained by using the above-mentioned coordinates and anatomical landmarks for voxel placement on the set of sagittal and coronal 1H MR images. For metabolic fitting, spectra were acquired from equal voxels in spherical metabolite phantoms (0.1 mol/l metabolite, pH 7.2, 37°C). The point resolved spectroscopy (PRESS) sequence preceded by water suppression (3 Gauss chemical shift selective pulses of 25.6 ms duration) was used throughout. For one metabolite spectrum, eight subspectra of 16 phase cycled scans each were recorded with a repetition time of 3 s. The eight metabolite subspectra were corrected for eddy currents using water-unsuppressed spectra ( $n=8$ ), automatically corrected for frequency and phase shifts, and added together to give 128 averages. Spectral quantification was carried out using a time domain–frequency domain fitting procedure (Schubert *et al.*, 2004) that includes phantom basis spectra and prior knowledge and involves background estimation by regularization. In the present spectra, the amplitudes of total choline, total creatine, *N*-acetylaspartate (NAA), glutamate and glutamine resonances were fitted. Residual, minor contributions by macromolecules are accommodated in the baseline by the fitting procedure. Using an echo time of 80 ms, the applied methodology has



**Fig. 1** Voxel positions shown on typical brain MDEFT images; upper panel: left dorsolateral prefrontal cortex; lower panel: visual cortex.


been shown to yield maximum selectivity for the glutamate C4 resonance over glutamine and gamma-aminobutyric acid (Schubert *et al.*, 2004). Mean uncertainties [corresponding to Cramér–Rao lower bounds with added uncertainties from background modelling (Elster *et al.*, 2005)] for the fitting of glutamate as small as 12.1% for the DLPFC voxel and 8.9% for the VC voxel were obtained. The fitted glutamate amplitudes were corrected for different coil loading by the phantoms and the individual subject's head (principle of reciprocity) and for relaxation effects using relaxation times measured previously (Schubert *et al.*, 2004), assuming equal glutamate relaxation times in the investigated regions. Glutamate concentrations were corrected for the cerebrospinal fluid in the voxels studied by using the cerebrospinal fluid fractions obtained by segmenting the T1-weighted images with SPM2 (Wellcome Trust Centre for Neuroimaging, London, UK; <http://www.fil.ion.ucl.ac.uk/spm/spm2.html>).

### Statistical analyses


In this study, we focused on extraversion as the principal personality trait of interest. Median-split was used to create two groups of subjects with high, i.e. extraverts, and low, i.e. introverts, scores of extraversion on the NEO-FFI, respectively. To ensure the differentiation of the two groups, we confirmed our selection with normative data from a large German sample (Körner *et al.*, 2008), which showed that subjects in our two groups scored indeed more than 1 s.d. above or below the average score for the normative group. Statistical calculations were carried out as indicated in the 'Results' section using SPSS for Windows (Release 16.0). Group differences were evaluated with *t*-tests for independent samples. Within-group comparisons were performed using paired *t*-tests. Correlation analysis was performed with Pearson correlation. All tests were performed at a level of significance equal to 0.05.

### RESULTS

Extraverts had significantly lower glutamate concentration in the DLPFC than introverts ( $P=0.044$ ; Figure 2). In the extraverts, there were no differences in metabolite concentrations between the investigated brain regions. In the introverts, there was a trend for higher glutamate concentrations in DLPFC than in VC ( $P=0.054$ ; Figure 3). Concentrations of the other metabolites did not differ between introverts and extraverts. Correlations between metabolite concentrations and psychometric measures were performed separately for extraverts and introverts. In the introverts, glutamate concentration in DLPFC was associated with a lower state anxiety (STAI 1;  $r=-0.59$ ,  $P=0.044$ ). There was no significant correlation between DLPFC glutamate concentration and state anxiety in the extraverts ( $r=-0.18$ ,  $P=0.491$ ; Figure 4). The relationship between glutamate and anxiety was specific for the DLPFC. Glutamate concentration in VC



**Fig. 2** Bar diagrams show DLPFC glutamate concentrations in extraverts and introverts. A significant difference between the two groups was observed ( $P=0.044$ ). DLPFC = dorsolateral prefrontal cortex.




**Fig. 3** Bar diagrams show glutamate concentrations in DLPFC and the control region (VC) in extraverts and introverts. A difference in glutamate concentration between the two regions was observed in introverts ( $P=0.054$ ), but not in extraverts. DLPFC = dorsolateral prefrontal cortex; VC = visual cortex.

was not associated with state anxiety in introverts ( $r=0.053$ ) or in extraverts ( $r=-0.18$ ).

### DISCUSSION

This study has three major findings: (i) introverts have significantly higher DLPFC glutamate concentration than extraverts; (ii) introverts have higher glutamate concentration in DLPFC than in VC and (iii) in introverts there is a negative association between DLPFC glutamate concentration and self-reported measures of state anxiety.

Previous studies have emphasized the role of the prefrontal cortex in extraversion and reported a decreased bilateral cerebral blood flow (Mathew *et al.*, 1984; Stenberg *et al.*, 1990; Johnson *et al.*, 1999) as well as a relative regional thinning of the lateral prefrontal cortex (Wright *et al.*, 2006). More medial regions of the frontal lobe, on the other hand, show a positive correlation with extraversion (Deckersbach *et al.*, 2006; Simon *et al.* 2010; Brühl *et al.*, 2011).



**Fig. 4** Scatterplot of DLPFC glutamate concentrations and state anxiety (STAI 1) scores in extraverts and introverts. A significant negative correlation was observed in introverts, but not in extraverts.  $^*P < 0.005$ .

Also, cerebral blood flow and activation in the amygdala in response to positive stimuli correlate with extraversion (Canli *et al.*, 2002; Vaidya *et al.* 2007), and there is a positive association between extraversion and amygdala grey matter concentration (Omura *et al.*, 2005). An association with extraversion has thus been shown for both cortical and subcortical regions suggesting that neural systems associated with this specific personality trait are not confined to higher level executive brain regions, but are rather represented at all levels of neural processing (Canli 2004). Since there seems to be a dissociation between lateral prefrontal, medial prefrontal and amygdala structural and functional findings in extraversion, one might assume a differential role for these regions. Our finding of decreased DLPFC glutamate in extraverts might be the result of reduced DLPFC gray matter concentration and/or decreased activation reported in previous studies (Johnson *et al.*, 1999; Wright *et al.*, 2006) and is consistent with two of the most significant biological models of personality, both of which hold that extraverts have lower cortical activity, especially in the frontal lobes (Gray, 1972; Eysenck, 1986). Lower activity and reduced glutamatergic neurotransmission in the DLPFC, a crucial region within the inhibitory topdown control network may be associated with reduced strategic implementation of control over thoughts and actions (MacDonald *et al.*, 2000; Botvinick *et al.*, 2001). A reduced topdown control of DLPFC on subcortical regions in conjunction with increased grey matter concentration in the amygdala in extraverts might also explain why amygdala activation to positive stimuli was found to correlate positively with the degree of extraversion in a previous study (Canli *et al.*, 2002). Psychologically, these processes might be reflected in the behavioural characteristics of extraverts such as high approach and reward sensitivity and the propensity to experience a wide range of positive moods (McCrae and Costa, 2003). According to the above-mentioned biological

models of personality, higher DLPFC glutamate concentration in introverts due to higher activity in the behavioural inhibition system, namely the prefrontal cortex (Gray, 1972), reflects more engagement in cognitive processes, such as making plans for the future or problem solving (Gray, 1972; Eysenck, 1986). Unlike in extraverts, in the introverted subjects glutamate concentrations in DLPFC were higher than in the other cortical region investigated, i.e. the VC. This finding emphasizes the role of glutamatergic metabolism in the DLPFC as a key region for the implementation of cognitive control (MacDonald *et al.*, 2000). Even though variation of Glx (glutamate + glutamine) across different brain regions has been reported, these differences concerned increases along the midline and between tissue types (Doelken *et al.*, 2009) and can therefore not explain the differences in glutamate concentrations between cortical regions found in our study.

Previous MRS studies that investigated the role of glutamate for different personality traits focused mainly on the ACC, where a negative association between glutamate and sensation seeking (Gallinat *et al.*, 2007) could be shown. In patients with borderline personality disorder as well as in healthy controls, a positive correlation with self-reported impulsivity has been reported (Hoerst *et al.*, 2010). These differences in correlation patterns may be related to differences between these two personality traits. Even though impulsivity and sensation seeking are related constructs, current knowledge posits sensation seeking as one of several aspects of impulsivity (Meda *et al.*, 2009). On the other hand, the dimension of extraversion is central to the sensation seeking personality trait (Zuckerman, 2007). Our finding of lower DLPFC glutamate concentration in extraverts therefore confirms the results reported by Gallinat *et al.* (2007). In animals the response to novelty, a construct which can be associated with approach behaviour and high sensation seeking (Dellu *et al.*, 1992), has been extensively

investigated. Blockade of glutamate receptors such as the NMDA (*N*-methyl-*D*-aspartate)-receptor or AMPA (alpha-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid)-receptor has been found to increase novelty induced exploratory behaviour (Alvarez and Ruarte, 2001; Martínez *et al.*, 2002). In human studies glutamatergic dysfunction has been reported in psychiatric disorders marked by specific alterations in the production of positive emotional states and approach behaviour, such as depression or anxiety disorders (Mathew *et al.*, 2008; Sanacora *et al.*, 2008; Walter *et al.*, 2009; Hashimoto *et al.*, 2010). Not only have low trait levels of extraversion been linked to increased vulnerability to major depression and anxiety (Watson and Clark, 1997; Kotov *et al.*, 2010), but dysfunction of the DLPFC also appears to be a key feature of depression (Davidson *et al.*, 2003; Grimm *et al.*, 2008).

The finding of a negative association between DLPFC glutamate concentration and self-reported measures of state anxiety in introverts further confirms the hypothesized relationship between extraversion, DLPFC function and glutamatergic neurotransmission. Introverts show higher activity in regions of the behavioural inhibition system, such as the DLPFC (Gray, 1972), and have higher glutamate levels, which results in an increased top-down control of DLPFC on the amygdala as a critical structure in the fear network (Davis and Whalen, 2001). There might have been no association with trait anxiety due to the fact that an acute affective state is more strongly associated with extraversion (Gross and John, 1998). The relationship between DLPFC glutamate and state anxiety is specific for the DLPFC since there is no correlation with VC glutamate.

To our knowledge, only one study has investigated the association between extraversion and neurotransmitter concentrations (Goto *et al.*, 2010); and the authors reported a negative correlation with GABA/creatine ratios in the frontal lobes. In contrast to our examination, in this study the spectroscopic voxel was centred on the interhemispheric fissure.

There are several limitations to this study. The rather small sample size of this investigation has to be considered when interpreting the results. Furthermore, we confined the study to females within a limited age range since a previous study showed a significant age-related decline in glutamate in men (Chang *et al.*, 2009). Even though no gender differences have been reported for the extraversion dimension of the NEO-FFI (Chapman *et al.*, 2007), a replication of the results in a male sample is needed. Additionally, while we investigated a control region unrelated to extraversion, we cannot exclude lateralization effects due to the positioning of the DLPFC voxel in the left hemisphere. Several studies suggest that the left, but not the right DLPFC is specifically involved in approach-related positive affective states (Davidson and Irwin, 1999; Davidson *et al.*, 2002).

## CONCLUSIONS

The present study indicates for the first time increased glutamate levels in the DLPFC of introverts as compared with extraverts. The increased glutamate concentration was specific for the DLPFC and negatively associated with state anxiety. The hypothesis of altered top-down control of DLPFC due to reduced glutamate concentration as a function of extraversion should be tested in further studies in healthy subjects as well as in psychiatric disorders marked by reduced approach behaviour.

## Conflict of Interest

None declared.

## REFERENCES

- Alvarez, E.O., Ruarte, M.B. (2001). Role of glutamate receptors in the nucleus accumbens on behavioural responses to novel conflictive and non-conflictive environments in the rat. *Behavioural Brain Research*, 123, 143–53.
- Bagby, R.M., Ryder, A.G. (2000). Personality and the affective disorders: past efforts, current models, and future directions. *Curr Psychiatry Report*, 2(6), 465–72.
- Banich, M.T., Mackiewicz, K.L., Depue, B.E., Whitmer, A.J., Miller, G.A., Heller, W. (2009). Cognitive control mechanisms, emotion and memory: a neural perspective with implications for psychopathology. *Neuroscience & Biobehavioral Reviews*, 33, 613–30.
- Barber, A.D., Carter, C.S. (2005). Cognitive Control Involved in Overcoming Prepotent Response Tendencies and Switching Between Tasks. *Cerebral Cortex*, 15, 899–912.
- Borkenau, P., Ostendorf, F. (2008). NEO-Fünf-Faktoren-Inventar Hogrefe. Botvinick, M.M., Braver, T.S., Barch, D.M., Carter, C.S., Cohen, J.D. (2001). Conflict monitoring and cognitive control. *Psychological Review*, 108, 624–52.
- Brühl, A., Viebke, M.-C., Baumgartner, T., Kaffenberger, T., Herwig, U. (2011). Neural correlates of personality dimensions and affective measures during the anticipation of emotional stimuli. *Brain Imaging and Behavior*, 5(2), 1–11.
- Canli, T. (2004). Functional brain mapping of extraversion and neuroticism: learning from individual differences in emotion processing. *Journal of Personality*, 72, 1105–32.
- Canli, T., Amin, Z. (2002). Neuroimaging of emotion and personality: scientific evidence and ethical considerations. *Brain and Cognition*, 50, 414–31.
- Canli, T., Sivers, H., Whitfield, S.L., Gotlib, I.H., Gabrieli, J.D.E. (2002). Amygdala Response to Happy Faces as a Function of Extraversion. *Science*, 296, 2191.
- Canli, T., Zhao, Z., Desmond, J.E., Kang, E., Gross, J., Gabrieli, J.D. (2001). An fMRI study of personality influences on brain reactivity to emotional stimuli. *Behavioral Neuroscience*, 115, 33–42.
- Chang, L., Jiang, C.S., Ernst, T. (2009). Effects of age and sex on brain glutamate and other metabolites. *Magnetic Resonance Imaging*, 27, 142–5.
- Chapman, B.P., Lyness, J.M., Duberstein, P. (2007). Personality and medical illness burden among older adults in primary care. *Psychosomatic Medicine*, 69, 277–82.
- Costa, P.T. Jr, McCrae, R.R. (1989). *The NEO-PI/NEO-FFI manual supplement*. Odessa, FL: Psychological Assessment Resources.
- Costa, P.T. Jr, McCrae, R.R. (1992). Normal personality assessment in clinical practice: the NEO Personality Inventory. *Psychological Assessment*, 4, 5–13.
- Davidson, R.J. (2003). Darwin and the neural bases of emotion and affective style. *Annals of the New York Academy of Sciences*, 1000, 316–36.
- Davidson, R.J., Irwin, W. (1999). The functional neuroanatomy of emotion and affective style. *Trends in Cognitive Sciences*, 3, 11–21.

- Davidson, R.J., Lewis, D.A., Alloy, L.B., et al. (2002). Neural and behavioral substrates of mood and mood regulation. *Biological Psychiatry*, 52, 478–502.
- Davis, M., Whalen, P. (2001). The amygdala: vigilance and emotion. *Molecular Psychiatry*, 6, 13–34.
- Deckersbach, T., Miller, K.K., Klibanski, A., et al. (2006). Regional cerebral brain metabolism correlates of neuroticism and extraversion. *Depression and Anxiety*, 23, 133–8.
- Dellu, F., Mayo, W., Cherkaoui, J., Le Moal, M., Simon, H. (1992). A two-trial memory task with automated recording: study in young and aged rats. *Brain Res*, 588(1), 132–9.
- Depue, R., Lenzenweger, M. (2001). A neurobehavioral dimensional model. In: Livesley, W.J., editor. *Handbook of personality disorders*. New York: Guilford Press, pp. 136–76.
- Doelken, M.T., Mennecke, A., Stadlbauer, A., et al. (2009). Multi-voxel magnetic resonance spectroscopy of cerebral metabolites in healthy adults at 3 Tesla. *Academic Radiology*, 16, 1493–501.
- Elster, C., Schubert, F., Link, A., Walzel, M., Seifert, F., Rinneberg, H. (2005). Quantitative magnetic resonance spectroscopy: semi-parametric modeling and determination of uncertainties. *Magnetic Resonance in Medicine*, 53, 1288–96.
- Eysenck, H. (1986). Psychophysiology and personality: extraversion, neuroticism and psychoticism. In: Gale, A., Edwards, J.A., editors. *Physiological Correlates of Human Behavior: Individual Differences and Psychopathology*, Vol. 3, Orlando: Academic Press, pp. 13–30.
- Gallinat, J., Kunz, D., Lang, U.E., et al. (2007). Association between cerebral glutamate and human behaviour: the sensation seeking personality trait. *NeuroImage*, 34, 671–8.
- Goldberg, I.R. (1992). Development of markers for the Big-Five factor structure. *Psychological Assessment*, 4, 26.
- Goto, N., Yoshimura, R., Moriya, J., et al. (2010). Critical examination of a correlation between brain gamma-aminobutyric acid (GABA) concentrations and a personality trait of extroversion in healthy volunteers as measured by a 3 Tesla proton magnetic resonance spectroscopy study. *Psychiatry Research: Neuroimaging*, 182, 53–7.
- Gray, J. (1972). The psychophysiological basis of introversion-extraversion: a modification of Eysenck's theory. In: Nebylitsyn, V.D., Gray, J.A., editors. *The Biological Bases of Individual Behavior*. New York: Academic Press, pp. 182–205.
- Grimm, S., Beck, J., Schuepbach, D., et al. (2008). Imbalance between left and right dorsolateral prefrontal cortex in major depression is linked to negative emotional judgment: an fMRI study in severe major depressive disorder. *Biological Psychiatry*, 63, 369–76.
- Gross, J.J., John, O.P. (1998). Mapping the domain of expressivity: Multimethod evidence for a hierarchical model. *Journal of Personality and Social Psychology*, 74, 170–91.
- Hashimoto, K. (2010). The role of glutamate on the action of antidepressants. *Progress in Neuro-Psychopharmacology and Biological Psychiatry* 35(7), 1558–68.
- Hoerst, M., Weber-Fahr, W., Tunc-Skarka, N., et al. (2010). Correlation of glutamate levels in the anterior cingulate cortex with self-reported impulsivity in patients with borderline personality disorder and healthy controls. *Archives General Psychiatry*, 67, 946–54.
- Jang, K.L., McCrae, R.R., Angleitner, A., Riemann, R., Livesley, W.J. (1998). Heritability of facet-level traits in a cross-cultural twin sample: support for a hierarchical model of personality. *Journal of Personality and Social Psychology*, 74, 1556–65.
- Johnson, D.L., Wiebe, J.S., Gold, S.M., et al. (1999). Cerebral blood flow and personality: a positron emission tomography Study. *American Journal of Psychiatry*, 156, 252–7.
- Kelly, A.M.C., Hester, R., Murphy, K., Javitt, D.C., Foxe, J.J., Garavan, H. (2004). Prefrontal-subcortical dissociations underlying inhibitory control revealed by event-related fMRI. *European Journal of Neuroscience*, 19, 3105–12.
- Kim, H.J., Kim, J.E., Cho, G., et al. (2009). Associations between anterior cingulate cortex glutamate and [gamma]-aminobutyric acid concentrations and the harm avoidance temperament. *Neuroscience Letters*, 464, 103–7.
- Körner, A., Geyer, M., Roth, M., et al. (2008). Persönlichkeitsdiagnostik mit dem NEO-Fünf-Faktoren-Inventar: Die 30-Item-Kurzversion (NEO-FFI-30). *Psychotherapie, Psychosomatik, medizinische Psychologie*, 58, 238,245.
- Kotov, R., Gamez, W., Schmidt, F., Watson, D. (2010). Linking big personality traits to anxiety, depressive, and substance use disorders: a meta-analysis. *Psychological Bulletin*, 136, 768–821.
- Kumari, V., Pfyche, D.H., Williams, S.C.R., Gray, J.A. (2004). Personality predicts brain responses to cognitive demands. *The Journal of Neuroscience*, 24, 10636–41.
- Laux, L., Glanzmann, P., Schaffner, P., Spielberger, C.D. (1981). Das State-Trait-Angstinventar. Theoretische Grundlagen und Handanweisung. Beltz Test GmbH.
- Lavric, A., Pizzagalli, D.A., Forstmeier, S. (2004). When 'go' and 'nogo' are equally frequent: ERP components and cortical tomography. *European Journal of Neuroscience*, 20, 2483–8.
- Livesley, J.W. (2001). Conceptual and taxonomic issues. *Handbook of Personality Disorders: Theory, Research and Treatment*. New York: Guilford Press, pp. 3–33.
- Lucas, R.E., Diener, E. (2001). Understanding extraverts' enjoyment of social situations: the importance of pleasantness. *Journal of Personality and Social Psychology*, 81, 343–56.
- MacDonald, A.W., Cohen, J.D., Stenger, V.A., Carter, C.S. (2000). Dissociating the role of the dorsolateral prefrontal and anterior cingulate cortex in cognitive control. *Science*, 288, 1835–8.
- Martinez, G., Roperio, C., Funes, A., Flores, E., Blotta, C., Landa, A.I., Gargiulo, P.A. (2002). Effects of selective NMDA and non-NMDA blockade in the nucleus accumbens on the plus-maze test. *Physiology and Behavior*, 76, 219–24.
- Mathew, R.J., Weinman, M.L., Barr, D.L. (1984). Personality and regional cerebral blood flow. *British Journal of Psychiatry*, 144, 529–32.
- Mathew, S.J., Price, R.B., Charney, D.S. (2008). Recent advances in the neurobiology of anxiety disorders: implications for novel therapeutics. *American Journal of Medical Genetics Part C: Seminars in Medical Genetics*, 148C, 89–98.
- McCrae, R.R., Costa, P.T. Jr (1997). Personality trait structure as a human universal. *American Psychologist*, 52, 509–16.
- McCrae, R.R., Costa, P.T. (2003). Personality in adulthood, a five-factor theory perspective. New York: Guilford Press.
- McCrae, R.R., Costa, P.T. (1999). A five-factor theory of personality. In: Pervin, L.A., John, O.P., editors. *Handbook of personality: Theory and research*. New York: Guilford Press, pp. 139–53.
- Meda, S.A., Stevens, M.C., Potenza, M.N., et al. (2009). Investigating the behavioral and self-report constructs of impulsivity domains using principal component analysis. *Behavioural Pharmacology*, 20, 390–9.
- Montag, C., Schubert, F., Heinz, A., Gallinat, J. (2008). Prefrontal Cortex Glutamate Correlates with Mental Perspective-Taking. *PLoS ONE*, 3, e3890.
- Omura, K., Todd Constable, R., Canli, T. (2005). Amygdala gray matter concentration is associated with extraversion and neuroticism. *Neuroreport*, 16, 1905–8.
- Rajkowska, G., Goldman-Rakic, P.S. (1995). Cytoarchitectonic definition of prefrontal areas in the normal human cortex: II. Variability in location of areas 9 and 46 and relationship to the Talairach Coordinate System. *Cereb Cortex*, 5(4), 323–37.
- Sanacora, G., Zarate, C.A., Krystal, J.H., Manji, H.K. (2008). Targeting the glutamatergic system to develop novel, improved therapeutics for mood disorders. *Nature Review Drug Discovery*, 7, 426–37.
- Santor, D.A., Bagby, R.M., Joffe, R.T. (1997). Evaluating stability and change in personality and depression. *Journal of Personality and Social Psychology*, 73, 1354–62.
- Schubert, F., Gallinat, J., Seifert, F., Rinneberg, H. (2004). Glutamate concentrations in human brain using single voxel proton magnetic resonance spectroscopy at 3 Tesla. *NeuroImage*, 21, 1762–71.

- Sheehan, D.V., Lecrubier, Y., Sheehan, K.H., et al. (1998). The Mini-International Neuropsychiatric Interview (M.I.N.I.): the development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *Journal of Clinical Psychiatry*, 59, 22–33.
- Simon, J.J., Walther, S., Fiebach, C.J., et al. (2010). Neural reward processing is modulated by approach- and avoidance-related personality traits. *NeuroImage*, 49, 1868–74.
- Stenberg, G., Risberg, J., Warkentin, S., Rosen, I. (1990). Regional patterns of cortical blood flow distinguish extraverts from introverts. *Personality and Individual Differences*, 11, 663–73.
- Sugiura, M., Kawashima, R., Nakagawa, M., et al. (2000). Correlation between Human Personality and Neural Activity in Cerebral Cortex. *NeuroImage*, 11, 541–6.
- Vaidya, J.G., Paradiso, S., Andreasen, N.C., Johnson, D.L., Boles Ponto, L.L., Hichwa, R.D. (2007). Correlation between extraversion and regional cerebral blood flow in response to olfactory stimuli. *American Journal of Psychiatry*, 164, 339–41.
- Walter, M., Henning, A., Grimm, S., et al. (2009). The relationship between aberrant neuronal activation in the pregenual anterior cingulate, altered glutamatergic metabolism, and anhedonia in major depression. *Archives of General Psychiatry*, 66, 478–86.
- Watson, D., Clark, L. (1997). Measurement and mismeasurement of mood: recurrent and emergent issues. *Journal of Personality Assessment*, 68, 267–96.
- Wright, C.I., Williams, D., Feczko, E., et al. (2006). Neuroanatomical correlates of extraversion and neuroticism. *Cerebral Cortex*, 16, 1809–19.
- Zuckerman, M. (2007). The sensation seeking scale V (SSS-V): still reliable and valid. *Personality and Individual Differences*, 43, 1303–5.